

- KITCHENER, D.J. & CHAPMAN, A. 1978. Mammals of Dongolocking Nature Reserve. In *Biological Survey of the Western Australian Wheatbelt, Part 5. Rec. West. Aust. Mus. Suppl. 6*: 53-58.
- MILTON, D.A. 1980. A Comparison of three techniques used in reptile survey of the Conondale Ranges. *Vic. Nat.* 97: 26-31.
- MORRIS, K.D. & KITCHENER, D.J. 1979. Mammals of Yornaning Nature Reserve. In *Biological Survey of the Western Australian Wheatbelt, Part 7. Rec. West. Aust. Mus. Suppl. 8*: 29-33.
- SERVENTY, D.L. & WHITTELL, H.M. 1976. *Birds of Western Australia*. Perth: University of Western Australia.
- STORR, G.M. 1975. The genus *Hemiergis* (Lacertilia, Scincidae) in Western Australia. *Rec. West. Aust. Mus.* 3(4): 251-260.
- STORR, G.M. 1976. The genus *Menetia* (Lacertilia Scincidae) in Western Australia. *Rec. West. Aust. Mus.* 4(2): 189-200.
- STORR, G.M. 1978. Whip snakes (*Demansia*, Elapidae) of Western Australia. *Rec. West. Aust. Mus.* 6(3): 287-301.
- STORR, G.M. & JOHNSTONE, R.E. 1979. *Field Guide to the Birds of Western Australia*. Perth: Western Australian Museum.
- Worsley Environmental Review and Management Programme. 1978. Perth: Maunsell and Partners Pty. Ltd. pp. 5.11-5.14.
- Worsley Alumina. 1981. Flora and Fauna Studies, Phase One. Worsley Alumina Pty. Ltd. and Dames and Moore, Perth.

FROM FIELD AND STUDY

Spotted harrier: a new record for Rottnest Island — On June 15, 1983, we saw an adult Spotted harrier *Circus assimilis* drifting across heathland near the Cathedral Rocks lookout at the west end of the island. The bird moved westward, following the contours of the dunes and heathlands in characteristic flight, and was last seen flying eastward towards Radar Reef.

On the following day Mr. Ian White, the resident lighthouse keeper, reported that he had seen what he believed was a Spotted harrier in the vicinity of the lighthouse on Wadjemup Hill. The single bird was seen 6 times on the following dates: 22,23 January 1983; some time before and including 28 March 1983; 11 May 1983 and 18 May 1983.

Serventy (1976) considers the Spotted harrier to be essentially an arid-country bird that periodically strays onto the coastal plain of south Western Australia. Milhinch and Ritcher (1979) found this species breeding in the Northam district in 1978, but the birds appeared to have deserted the area by January 1979.

REFERENCES

- MILHINCH, A.L. and RICHTER, C.M. 1979. *Nesting Spotted harrier near Northam*. W.A. Naturalist, 14: 130.
- SERVENTY, D.L. and WHITTELL, H.M. 1976. *Birds of Western Australia*, 5th ed. University of Western Australia Press: Perth.
- C.P. de REBEIRA and D.A. SAUNDERS, CSIRO, Division of Wildlife & Rangelands Research.